



ORIGINAL RESEARCH ARTICLE

ORAL MEDICATION ADMINISTRATION PRACTICE OF NURSES IN CHITWAN MEDICAL COLLEGE L Rajbanshi ^{1*}

¹ College of Nursing, Chitwan Medical College, Bharatpur, Chitwan

*Correspondence to: Ms. Laxmi Rajbanshi, Associate Professor & Matron, College of Nursing, Chitwan Medical College. Email: laxmiraj3@yahoo.com

ABSTRACT

Administering medication is high risk activity because one third of the errors occur during the nurse administration phase. Objective of the study was to find out the nurses practice of administering medicine to the patients. For that on duty nursing supervisors were observed the activities of nurses during administering medicines to the patients at 6 pm, 10 pm and 6 am of following day. Out of 52 nurses, only 5.76% nurses asked patients about the history of allergy and 86.53% asked patients to state their name for identification, 55.76% of nurses compared label of drug with cardex while withdrawing from the stock, 38.46% while pouring and 40.38% before replacing the drug on stock., 13.46% nurse explained purpose of drug to patient, 94.23% were check expiry date of drug, 46.15% stayed with patient till swallowing of drug. This study revealed, 68.42% nurses used measuring cup for accurate measurement of the liquid drug. Similarly, transparency of the syrup was checked by the 50.0% of nurses and 73.60 % of nurses shook the drug(suspension) before pouring on cup. As nurses were poorly adhere to the medication administration procedure, development of policy and procedure with ongoing education and monitoring of practice on safe medication is recommended.

Key words: Administration, Medication, Nurse, Practice.

DOI: <http://dx.doi.org/10.3126/jcmc.v6i2.16680>

INTRODUCTION

The administration of a medicine is a common but important clinical procedure. Administering medication is high risk activity because one third of the errors occur during the nurse administration phase.¹ It is the manner in which a medicine is administered that will determine to some extent whether or not the patient gains any clinical benefit, and whether they suffer any adverse effect from their medicines.² However Nursing Interventions Classification (NIC), defined medication administration as preparing, giving and evaluating effectiveness of prescription and nonprescription medications.³

Medication administration is complex multistep process that encompasses, prescribing, transcribing, dispensing, administering drug and monitoring responses.⁴ It is more than a simple psychomotor task. It reflects a complex interaction of a large number of specific decisions and actions⁵ and involves skillful technique and consideration of the

patients' development, health status and safety. The nurse administering medication needs knowledge about drugs including drugs name, preparation, classifications and adverse effects and physiological factors that affect drug actions.⁶ Administering medication safely requires an understanding of legal aspects of health care, pharmacology, pharmacokinetics, the life science, human anatomy and mathematics⁷ because a study revealed, the main causes of medication administration errors were lack of calculation, competency and poor adherence to protocol, distractions and time pressures.⁸

Medicine can be administered by the different routes such as parental, oral, topical and pulmonary. Among all, oral route is the most frequently used route of drug administration and is the most convenient and economic route. In oral route, patient's needs to swallow drug and drug given orally are intended for

absorption in the stomach and small intestine.⁶

In hospital, nurses are primarily responsible for providing medicine to the patient. Administering medication is a high risk activity because one third of the error occurs during the nurse administration phase¹ as once the medicine has been given, there is no way to intercept it. Therefore, errors in medication administration are more crucial than errors in other stages of the medication process (prescribing, transcribing, dispensing and monitoring) as 84% of these errors go un-intercepted⁹ and it may cause serious harm to the patient.¹⁰ Thus, safe administration of medication is very important to the nurse, doctor, administrator, patient and the public and the entire health care system.⁵ Administering medicines to the patients according to the defined procedure and policy reduces the risk of error and assures the patient's safety. Therefore, this study aims at finding out the nurses' practice of administering medicine to the patients by oral route.

OBJECTIVE

To find out the oral medication administration practice of nurses in CMC Teaching Hospital.

METHODOLOGY

This observational study was carried out at Chitwan Medical College (CMC) on Baishakha, 2069.

Observation check list was used to collect the data. Night nursing supervisors were mobilized for observing the activities of nurses while administering medication at 6 pm, 10 pm and 6 am of the following day. Nurses working in Surgery, Medicine, Orthopaedics and Gynaecology wards were included in the study. Total 52 nurses' practice was observed. Frequency and percentage were used to analyze the data.

RESULT

Out of 52 nurses, 17.30% of nurses checked the patient's ability to swallow medicine, 11.53% asked the history of vomiting/diarrhea or NPO and only 5.76% of nurses asked the history of allergy and none of them checked the bowel sound before administering the drug. Similarly, during administration of the drug, hand washing was performed by 48.07% of nurses and 86.53% asked the patients to state their name for identification (table 1). While administering liquid medicine, 73.60% of nurses shook the medicine before pouring it into the measuring cup, 50% were observed to check its transparency and 68.42% used a calibrated medicine cup for measurement. Furthermore, after administering medicines to the patients, 94.23% immediately signed on the Cardex; however, the patient's response to the medicine (after 30 minutes of administration) was assessed by 11.53% of nurses and hand hygiene was performed by 44.23% of nurses.

Table 1 Steps Performed by Nurses during Administration of Medicine (n=52)

Steps	Frequency	Percentage
Perform hand washing	25	48.07
Prepare medication of one patient at a time	50	96.15
Compare label of drug with Cardex:		
• While withdrawing from stock (drawer)	29	55.76
• Before pouring on container	20	38.46
• Before replacing to stock (drawer)	21	40.38
Check expiry date of drug	49	94.23
Perform dose calculation	41	78.84
Pour tablet in paper bag / medicine cup	22	42.30
Ask patient to state his/her name	45	86.53
Explain purpose of drug to patient	07	13.46
Place patient in sitting/side lying	35	67.30
Give one medicine at a time	41	78.84
Stay with patient until swallowing of drug	24	46.15

DISCUSSION

The distribution and administration of medication represents a major duty of hospital-based nurses.

This study revealed, out of 52, only 5.76% nurses asked patients about the history of allergy to that drug. The reason behind it was either, they think that the drug couldn't cause allergy or they may have asked during previous administration. However, a study conducted in Public Hospital of Soudi Arabia found 22% of nurses reported that the drug was given to known allergy patients.⁵ Only 11.53% nurses asked patients about the vomiting and diarrhea before administering the drug. None of them auscultated the bowel sound, as it is crucial for the patients with major abdominal surgery.

The "rights" of medication administration include right patient, right drug, right time, right route, and right dose.¹¹ These rights are critical for nurses. Accurate demographic information (the "right patient") is the first of the "five rights" whereas in this study 86.53% of nurses asked patients to state his/her name. This study found, 55.76% of nurses compared label of drug with cardex while withdrawing from the stock, 38.46% while pouring and 40.38% before replacing the drug on stock. These three checks are essential to ensure the right drug of medication administration because administration of wrong drug to a patient is second most common error among the medication errors.¹²

This study revealed, 68.42% nurses used measuring cup for accurate measurement of the liquid drug, and 60.05% held it at eye level while pouring. Similarly, transparency of the syrup was checked by the 50.0% of nurses and 73.60 % of nurses shook the drug (suspension) before pouring on cup, because failure to shake leads to low dose or overdose depending upon the drug product and elapsed time since the last use.

Hand hygiene is most crucial step of medication administration. This study showed only 48.07 % nurse performed hand hygiene before administration of drug and 44.23% after the procedure. Furthermore 42.30% nurses used paper bag or medicine cup for pouring medicine and 46.15% nurses stayed with patient until they swallowed medicine. After administration of drug 11.53% nurses assessed the patient again for monitoring the effect of drug administered.

Patients with impaired swallowing are always at risk of aspiration⁷ though it was assessed by only 17.30% of nurses and similarly, patient having medicines in an upright and side lying position, helps prevent the patient from choking and aspiration⁶ however, 67.30% of nurses placed patient in sitting/side lying position during administration.

CONCLUSION

This study concludes that nurses are poorly adhere to procedural protocol of the medication administration, which increase the risk of medication errors by the nurse and compromise the patient safety. Thus, for the prevention of errors, organization needs to develop medication administration policy, procedure and protocol along with provision of continue orientation, and supervision to nursing staff. Further study can be done on large setting and on prevalence of medication error and its associated factors.

ACKNOWLEDGEMENT

The authors would like to acknowledge the invaluable assistance of all Nursing Officers CMC Teaching Hospital for data collection, Dr. Anee in the preparation of this manuscript,

REFERENCES

1. Cloete L. Reducing medication errors in nursing practice. *Cancer Nursing Practice* 2015;14(1):29-36.
2. Bates D, Cullen D, Cooper J, et al. Systems analysis of adverse drug events. *JAMA* 1995;274:1599-1603.
3. Bulechek G, Butcher H, Dochterman J. (Eds.) (2008). *Nursing Interventions Classification (NIC)* (5th ed.) St. Louis: Mosby/Elsevier.
4. Anderson P, Townsend T. Medication Errors: Dont let them happen to you. *American Nurse Today*. March 2010;5(3): 23-27. .o
5. Ali Al-Youssifi S, Mohamed LK, Salim Mohamed NS. Nurses' Experiences toward Perception of Medication Administration Errors Reporting. *IOSR Journal of Nursing and Health Science*. May-Jun 2013;1(4):56-70. www.iosrjournals.org retrieved on 30 May 2015

6. Taylor CR, Lillis C, LeMone P, Lyne P. Fundamental of nursing (7th Ed):The art and science of care .New Delhi, Wolters Kluwer Pvt.Ltd , India. 2012; 740
7. Potter PA, Perry AG. Fundamentals of Nursing (7th ed), Elsevier India Mosbey; 2009: pp 719 .
8. Jones SW. Reducing medication administration errors in nursing practice. Nursing Standard 2009;23(50):40-46. <http://dx.doi.org/10.7748/ns2009.08.23.50.40.c7213>
9. Leape LL, Bates DW, Cullen DJ, et al. Systems analysis of adverse drug events. JAMA 1995; 274(1):35-43.
10. Kale A, Keohane CA, Maviglia S, Gandhi TK, Poon EG. Adverse drug events caused by serious medication administration errors. BMJ Qual Saf 2012. doi: 10.1136/bmjqs-2012-000946 Pp
11. Copping C. Preventing and reporting drug administration error. Nursing Times. August 2005;101(33):32-34.
12. Hughes RG, Blegen MA. Patient Safety and Quality: An Evidence-Based Handbook for Nurses. Available on http://www.ncbi.nlm.gov/books/nbk2656_nbk_2656_dtls on with conservative management as revealed by this study. Moreover, the parents feel happy when their infants' nasolacrimal duct obstruction gets resolved without the need for a surgical procedure.